



TECHNICAL PROTOCOL
FOR
FRT-PGK-gb2-hygro-FRT
FRT flanked,
Pro- and Eukaryotic
Hygromycin Selection Cassette
(A010)

Gene Bridges GmbH
Im Neuenheimer Feld 584
69120 Heidelberg, Germany
Tel + 49 (0)6221 13708 11
Fax + 49 (0)6221 13708 29
Email: contact@genebridges.com
www.genebridges.com

CONTENTS

1 Eppendorf tubes + manual

1. FRT-PGK-gb2-hygro-FRT: PCR template (50 ng/μl, 20μl)
2. This manual

Store tube at -20°C

Please read

The products listed in this manual are for research purposes only. They are not designed for diagnostic or therapeutic use in humans, animals or plants. The Red[®]/ET[®] recombination technology is the intellectual property of Gene Bridges GmbH.

Conditions of use

3.1 Purchaser will not manufacture, copy, reproduce, transmit, distribute, sell, lease, transfer, or improve upon the MATERIALS without prior written consent from GENE BRIDGES.

3.2 All MATERIALS relating Technologies shall be purchased from GENE BRIDGES or its authorized distributors. Use of any of the stated products from a source other than GENE BRIDGES will exempt GENE BRIDGES from any and all liabilities and warranties.

3.3 All MATERIALS purchased by research organizations, universities and other non-profit organizations may not be used for any commercial purpose. These MATERIALS are to be used for research purposes only. The MATERIALS may not be used to provide a commercial or non-commercial service, of any kind.

3.4 A purchase of MATERIALS by a private consumer is neither intended nor permitted.

Short Description:

“FRT-PGK-gb2-hygro-FRT” template is designed to allow hygromycin selection in prokaryotic and eukaryotic cells.

The FRT-PGK-gb2-hygro-FRT template encodes the hygromycin resistance gene which combines a prokaryotic promoter (gb2) for expression in *E.coli* with a eukaryotic promoter (PGK) for expression in mammalian cells.

The prokaryotic promoter gb2 is a slightly modified version of the Em7 promoter; it mediates higher transcription efficiency than the normally used Tn5 promoter. The promoter of the mouse Phosphoglucokinase gene (PGK) is used as eukaryotic promoter. A synthetic polyadenylation signal terminates the hygromycin expression. The cassette is flanked by FRT sites for later excision by FLP-recombinase.

Using the provided PCR template one can easily create a FRT-PGK-gb2-hygro-FRT cassette flanked by any restriction sites to clone the cassette into the vector of choice. The restriction sites can be introduced by adding the corresponding sequence in the PCR primer. The template can easily be used to generate targeting constructs mediated by a single Red/ET Recombination step.

The “FRT-PGK-gb2-hygro-FRT template” is not linear but plasmid based (3643bp in size). Due to its R6K origin it can not replicate in most of the *E. coli* strains. The PCR product can therefore be used directly for downstream applications without any further purification.

We recommend growing the recombined cells between 4 and 24h under non-selective conditions before streaking the culture on plates conditioned with 100 µg/ml hygromycin plus the antibiotics for the selective marker of the target vector (e.g. 15 µg/ml chloramphenicol for BAC clones).

At least 20 PCR reactions can be performed using 1µl per reaction as template.

Map: FRT-PGK-gb2-hygro-FRT cassette

